

EOLOGIC MAP OF SHEEP CANYON QUADRANGLE, WYOMING

by

Robert E. Ladd

1986

EXPLANATION

Qt₁
Qt₂
Qt₂
Qt₃

Terrace Levels
Qt Terraces along the Bighorn River and the Greybull River. Qt₁ is lowest surface; Qt₆ is highest.

 Qt 4

 Qt 5

 Qt 6

Qls

Landslide Material

Qal Unconsolidated deposits of alluvium along stream valleys at or near present stream levels.

Blocks of bedrock or loose slope debris.

Qaf
Alluvial Fan
Unconsolidated fan-shaped deposits of alluvium.

Colluvium

Heterogeneous deposits of rock detritus.

Willwood Formation

Variegated reddish-brown to olive-gray siltstone interbedded with lenticular, light gray,
ledge-forming sandstone. Thickness 2,500 feet.

Tfu

Fort Union Formation

Upper part is olive-gray siltstone, gray shale, thin brown to black coaly layers, and some lenticular sandstones. Lower part is lenticular, yellowish-gray sandstone layers interbedded with olive-gray siltstone and thin-bedded

Lance Formation

Light gray sandstone interbedded with gray to brown carbonaceous shale. Fossil bone fragments and round sandstone concretions in lower part. Thickness 800 feet.

carbonaceous shale. Thickness 3,750 feet.

Km

Meeteetse Formation

Yellowish-gray sandstone interbedded with brown to black carbonaceous shale and coal. Some thin bentonite layers. Thickness 850 feet.

Mesaverde Formation
Upper thick-bedded or massive, cliff-forming, light gray sandstone interbedded with olive-gray shale. Contains small iron concretions and fossil

wood. Lower thick-bedded, gray shale with thin-bedded, yellowish-gray, ledge-forming sand-stone. Thickness 1,350 feet.

Cody Shale

Dark gray to olive-brown, fissile shale. Contains layers of isolated septarian concretions.
Thickness 2,400 feet.

Kf
Frontier Formation
Upper Torchlight Sandstone Member is light

Frontier Formation
Upper Torchlight Sandstone Member is light gray sandstone with andesite, chert, and quartzite pebbles near top. Middle part is dark gray shale, friable, light gray sandstone and thin bentonite layers. Basal Peay Sandstone Member is thick-bedded, gray sandstone.

Mowry Shale
Upper 400 feet is siliceous, gray shale with thin, indurated sandstones near top. Contains fish scales, bentonite layers, and cone-in-cone structures near base. Lower 230 feet is soft.

with the Mowry to reflect the most recent accepted nomenclature. Thickness 630 feet.

Thermopolis Shale

Black, fissile shale and olive-gray to brownish-black shale with thin, olive-gray sandstone layers. Contains dahllite concretions. Upper 20 feet is Muddy Sandstone Member (Kmd) which is light gray, bentonitic sandstone with

dark gray, fissil shale and thin bentonite layers. Lower part was mapped as the Shell Creek Shale by the author, but has been included

Cloverly Formation
Variegated red, grayish-purple, and yellowish-brown mudstone and shale interbedded with thin-bedded, sandstone and siltstone near top. Upper 100 feet is mapped as Sykes Mountain Member (Ksm) which is yellowish-brown shale with thin sandstone layers overlying light orange sandstone interbedded with

thin layers of black shale.

Morrison Formation

Variegated gray-green mudstone and siltstone.

Yellowish-gray, friable sandstone at base.

Thickness 300 feet.

light gray shale and siltsone. Thickness 250

Sundance Formation

Upper Sundance is gray-green shale with interbedded glauconitic sandstone and fossiliferous limestone. Contains zones of Pachyteuthis densus, Ostrea engelmanni, and Camptonectes bellistriatus. Lower Sundance is gray-green, glauconitic siltstone and shale. Contains a zone of Gryphaea nebrascensis. Distinctive limestone layers occur near base and middle. Thickness

Gypsum Spring Formation

Upper part is red siltstone. Middle part is interbedded limestone and red shale. Lower part is massive gypsum. Thickness 220 feet.

Reddish-brown, calcareous siltstone with some layers of fine-grained sandstone. This formation was mapped by the author as the Red Peak Formation in the Chugwater Group. Its name has been changed to match adjoining maps and to avoid confusion. Thickness 570 feet.

Chugwater Formation

Dinwoody Formation

Bluish-gray siltstone and mudstone with a boxwork of thin gypsum stringers. The Dinwoody and underlying Phosphoria could just as well be called the Goose Egg Formation here, since there is an intertonguing relationship in this area. Thickness 55 feet.

Phosphoria Formation
Upper part is gray limestone or coarse limestone breccia with some interbedded chert and red shale layers. Lower part is thick-bedded gypsum with interbedded red siltstone or stromatolitic limestone. Basal part is distinctive reddish-brown siltstone. The Phosphoria and overlying Dinwoody could just as well be called the Goose Egg. Formation here, since there is an intertonging relationship in this area. Thickness 250 feet.

Pt
Upper part is massive to thick-bedded, light gray sandstone with thin shale and chert layers.
Lower part is interbedded, thin sandstone, limestone, siltstone, and chert layers. Thickness 95 feet.

Amsden Formation

Upper part is thin, purple shale, siltstone, and gray limestone layers overlying a distinctive cherty limestone. Lower part is red siltstone and shale which sometimes contains pisolitic layers. Basal Darwin Sandstone Member is light gray sandstone. Thickness 120 feet.

Madison Limestone
Upper part is limestone and limestone breccia
and contains caves. Lower part is massive,
gray to olive-gray limestone. Thickness 760

Formation Contact

Dashed where approximately located; dotted where concealed.

Trace of axial plane and direction of plunge.

Dashed where approximately located.

Clastic Dike
Dashed where approximately located.

Strike and dip of beds